

Comment on ‘Two-body random ensembles: from nuclear spectra to random polynomials’

R. Bijker^a and A. Frank^{a,b}

^a*Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México,
Apartado Postal 70-543, 04510 México, D.F., México*

^b*Centro de Ciencias Físicas, Universidad Nacional Autónoma de México,
Apartado Postal 139-B, Cuernavaca, Morelos, México*

An investigation of the spectroscopy of even-even nuclei with random one- and two-body interactions has led to the surprising result that the ground state has $J^P = 0^+$ in typically 60 – 70 % of the cases. This holds both for the nuclear shell model [1,2] and the interacting boson model [3]. To understand this phenomenon it was suggested to study the distribution of the lowest eigenvalues [4]. To this end, a technique was developed in [5] for tridiagonal Hamiltonian matrices, such as for the one- and two-body interactions in the vibron model. It relies on the property that for these matrices it is possible to obtain an estimate for the lowest eigenvalue as the minimum of a function [6].

The purpose of this comment is to point out that for the vibron model the appropriate function is not a quadratic polynomial as used in [5], but a more general form which in the vibrational basis to leading order in $1/N$, is given by

$$f(z, j) = az^2 + bz + c + dj^2 + e\sqrt{(1-z)^2(z^2 - j^2)}.$$

Here $z = n_p/N$ and $j = J/N$ [5]. In the absence of off-diagonal terms ($e = 0$), the spectrum is that of a three-dimensional oscillator characterized by $n_p = 0, 1, \dots, N$ and $J = n_p, n_p - 2, \dots, 1$ (n_p odd) or 0 (n_p even). In this case, the function $f(z, j)$ reduces to a quadratic polynomial, and the results of [5] can be used to obtain the percentages for ground states with $J = 0$, $J = 1$ and $J = N$ to be 64.6, 5.2 and 19.8 %, respectively, in close agreement with the exact values of 64.4, 4.1 and 24.2 % obtained for $N = 20$ and 100000 runs. In the case of only off-diagonal interactions ($a = b = c = d = 0$), the ground state always has $J = 0$.

In the general case, both diagonal and off-diagonal terms are present. The off-diagonal term introduces a deformation in the system, which gives rise to the occurrence of vibrational bands with $J = 0, 1, 2, \dots, J_{\max}$. As a consequence, the $J = 1$ ground states tend to disappear in favor of an enhancement of the $J = 0$ ground states. This effect explains qualitatively the change in the $J = 0$, $J = 1$ and $J = N$ ground states from 64.4, 4.1 and 24.2 to 72.1, 0.5 and 23.7 %, respectively.

In order to carry out a more quantitative analysis, we have developed a simple method to analyze arbitrary functions $f(z, j)$ by sampling the two-dimensional space (z, j) . As an example, in Figure 1 we show the percentages of ground states with $J = 0$ as a function of

$x = \sigma_1/(\sigma_0 + \sigma_1)$, where σ_i denotes the width of the Gaussian distributions for the diagonal ($i = 0$) and off-diagonal ($i = 1$) matrix elements. The case of equal widths (discussed in the previous paragraph) corresponds to $x = 0.5$. Figure 1 shows a good agreement between the values obtained from an analysis of $f(z, j)$ (dashed line) and the exact results (solid line). The discrepancies observed for the quadratic polynomials of [5] (dotted line) are due to a different treatment of the off-diagonal matrix element.

In conclusion, the approach proposed in [5] provides an interesting method to analyze random interactions in terms of random polynomials which is especially suited for tridiagonal Hamiltonian matrices. The advantage of this method is that it foregoes the diagonalization of thousands of matrices. In this comment we have presented a careful analysis of the dominance of $J^P = 0^+$ ground states in the vibron model, and found good agreement with the exact calculations.

For cases of interest to nuclear physics, such as the IBM or the shell model, the Hamiltonian matrix has a more complicated form. Although it can be brought to tridiagonal form, the lack of an analytic form for the function f leads to complications comparable to those of bringing it to diagonal form right away. For these cases the method becomes difficult to apply (other than by analogy). Many questions remain open to explain the robust features observed in randomly interacting many-body systems.

This work was supported in part by CONACyT under projects 32416-E and 32397-E, and by DPAGA under project IN106400.

-
- [1] C.W. Johnson, G.F. Bertsch and D.J. Dean, Phys. Rev. Lett. **80**, 2749 (1998).
 - [2] R. Bijker, A. Frank and S. Pittel, Phys. Rev. C **60**, 021302 (1999).
 - [3] R. Bijker and A. Frank, Phys. Rev. Lett. **84**, 420 (2000).
 - [4] R. Bijker and A. Frank, Phys. Rev. C **62**, 014303 (2000).
 - [5] D. Kusnezov, Phys. Rev. Lett. **85**, 3773 (2000).
 - [6] Lloyd C.L. Hollenberg and N.S. White, Phys. Rev. B **54**, 16309 (1996).




FIG. 1. Percentage of $J^P = 0^+$ ground states as a function of x (defined in text) for the exact results (solid), for $f(z, j)$ (dashed) and for the quadratic polynomial (dotted) [5].